

# PREPARATION OF HIGHLY ACTIVE H-PTH (1-34) AMIDE

**Publication number:** JP58096052 (A)

**Publication date:** 1983-06-07

**Inventor(s):** FUNAKOSHI SUSUMU

**Applicant(s):** TOYO JOZO KK

**Classification:**

**- international:** C07K14/575; A61K38/22; C07C67/00; C07C231/00; C07C231/02; C07C231/04; C07K14/00; C07K16/00; C07K14/435; A61K38/22; C07C67/00; C07C231/04; C07K14/00; C07K16/00; (IPC1-7): A61K37/24; C07C102/00; C07C103/52

**- European:**


**Application number:** JP19810193212 19811130

**Priority number(s):** JP19810193212 19811130

## Abstract of JP 58096052 (A)

**PURPOSE:**To prepare the titled high purity substance useful as a remedy for parathyroid insufficiency, by amidating a C-terminated phenylalanyl group, condensing protected amino acids in liquid phase in a specific sequence, eliminating the protective group of the N-terminal, and purifying the product by gel filtration, etc.

**CONSTITUTION:**The carboxyl group of the C-terminated phenylalanyl group is converted to amide group, and the protected amino acids and/or protected peptides are condensed by the liquid-phase synthesis in the order of formula (Ser is serine; Val is valine; Glu is glutamic acid; Ile is isoleucine; Gln is glutamine; Leu is leucine; Met is methionine; His is histidine; Asn is asparagine; Gly is glycine; Lys is lysine; Arg is arginine; Trp is tryptophane; Asp is aspartic acid residue), and the protective group of the N-terminal amino group is eliminated at the final stage of the condensation by acid decomposition. The product is separated and purified by the column chromatography using a gel filtration agent and an adsorbent.



---

Data supplied from the **esp@cenet** database — Worldwide